

# Some congruences on prime factors of class number of finite algebraic extensions $K/\mathbb{Q}$ , version 4.0

Roland Quême

2002 april 20

## Contents

|          |                                                                                                  |          |
|----------|--------------------------------------------------------------------------------------------------|----------|
| <b>1</b> | <b>On prime factors of class number of finite algebraic extensions <math>K/\mathbb{Q}</math></b> | <b>4</b> |
| 1.1      | Some definitions . . . . .                                                                       | 4        |
| 1.2      | Some results . . . . .                                                                           | 5        |
| <b>2</b> | <b>Geometry of Numbers point of view</b>                                                         | <b>7</b> |
| <b>3</b> | <b>Numerical examples</b>                                                                        | <b>8</b> |
| 3.1      | Cyclotomic number fields $\mathbb{Q}(\zeta_u)$ . . . . .                                         | 8        |
| 3.2      | Real class number $\mathbb{Q}(\zeta_l + \zeta_l^{-1})$ . . . . .                                 | 10       |
| 3.3      | Cubic fields $K/\mathbb{Q}$ cyclic and totally real. . . . .                                     | 11       |
| 3.4      | Totally real cyclic fields of prime conductor $< 100$ . . . . .                                  | 11       |
| 3.5      | Lehmer quintic cyclic field . . . . .                                                            | 11       |
| 3.6      | Decimic imaginary cyclic number fields with conductor between 9000 and 9500                      | 12       |
| 3.7      | Not abelian Galois solvable extensions $L/K$ . . . . .                                           | 12       |

# Abstract

Roland Quême  
 13 avenue du château d'eau  
 31490 Brax  
 France  
 2003 april 20  
 tel: 0561067020  
 mailto: roland.queme@free.fr  
 home page: <http://roland.queme.free.fr/>  
 \*\*\*\*\*

*Mathematical Subject Classification : 2000 M.S.C.*

Primary 11R29: Class Number  
 Secondary 12F10: Galois Theory  
 \*\*\*\*\*

- This paper is a contribution to the description of some congruences on the odd prime factors of the class number of the number fields.
- We say that a finite Galois extension  $L/K$  is Galois solvable if the Galois group  $Gal(L/K)$  is solvable. An example of the results obtained is:  
*Let  $L/\mathbb{Q}$  be a finite Galois solvable extension with  $[L : \mathbb{Q}] = N$ , where  $N > 1$  is odd. Let  $h(L)$  be the class number of  $L$ . Suppose that  $h(L) > 1$ . Let  $p$  be a prime dividing  $h(L)$ . Let  $r_p$  be the rank of the  $p$ -class group of  $L$ . Then  $p \times \prod_{i=1}^{r_p} (p^i - 1)$  and  $N$  are not coprime.*
- We give also in this paper a connection with Geometry of Numbers point of view. With an explicit geometric upper bound  $H_F$  of the class number  $h(F)$  for any field  $F$ , which is given in this paper:  
*Let  $L/\mathbb{Q}$  be a finite Galois extension with  $[L : \mathbb{Q}] = N$ . Let  $h(L)$  be the class number of  $L$ . Suppose that  $h(L) > 1$ . Suppose that  $N$  has odd prime divisors. Let  $q$  be an odd prime divisor of  $N$ . Then there exists a cyclic extension  $L/F$  with  $q = [L : F]$ . Suppose that  $p > H_F$  is a prime dividing  $h(L)$ . Let  $r_p$  be the rank of the  $p$ -class group of  $L$ . Then  $p \times (p^{r_p} - 1) \equiv 0 \pmod{q}$ .*
- The proofs are **elementary**. We give several verifications of results obtained for cyclic and abelian extensions from the tables in Washington[11], Schoof [10], Masley [8], Girstmair [4],

Jeannin [5] and the tables of cubic totally real number fields of the *ftp* server *megrez.math.u-bordeaux.fr*.

# 1 On prime factors of class number of finite algebraic extensions $K/\mathbb{Q}$

The so called *rank theorem*, see Masley [8], corollary 2.15 p. 305 is:

Suppose  $M/P$  is a cyclic extension of degree  $m$ . Let  $p$  be a prime which does not divide  $h(E)$ , class number of  $E$  for any field  $E$  with  $P \subset E \subset M$ ,  $E \neq M$ , and which does not divide  $m$ . If  $p \mid h(M)$  class number of  $M$  then the rank  $r_p$  of  $p$ -class group of  $M$  is a multiple of  $f$ , the order of  $p \bmod m$ .

In this subsection we use this theorem to get some congruences on prime factors of class number of some Galois extensions  $K/\mathbb{Q}$ .

## 1.1 Some definitions

- Let  $K/\mathbb{Q}$  be a finite algebraic extension. Let  $L/K$  be a finite Galois extension. We say that  $L/K$  is a *cyclic tower extension* if there is a tower of fields  $K = L_0 \subset L_1 \subset \cdots \subset L_n \subset L = L_{n+1}$  where  $L_{j+1}/L_j$ ,  $j = 0, \dots, n$ , is a cyclic extension. Observe that, without loss of generality, we can assume that  $[L_{j+1} : L_j]$  is a prime.
- We say that  $L/K$  is a Galois solvable extension if  $L/K$  is a Galois extension with a solvable Galois group  $Gal(L/K)$ .
- Let  $L/K$  be a finite extension. We say that  $L/K$  is a solvable extension if the smallest Galois extension  $M$  of  $K$  containing  $L$  is Galois solvable, see for instance Lang, [6] paragraph 7 p. 216. Classically, the Galois solvable extension  $M/K$  is a cyclic tower extension.
- Let  $L/\mathbb{Q}$  be a finite algebraic extension. Let  $N = [L : \mathbb{Q}]$ , where  $N = 2^{\alpha_0} \times N_1$  with  $N_1$  odd and where we assume that  $N_1 > 1$ .
- Let  $h(L)$  be the class number of the field  $L$ . In this paper, we are studying, for fields  $L$  with  $h(L) > 1$ , some congruences on the primes dividing  $h(L)$ .
- The class group  $C$  of  $L$  is an abelian group. Let  $p$  be a prime dividing  $h(L)$ . The  $p$ -subgroup  $C_p$  of  $C$  is an abelian group, direct sum of  $r_p$  cyclic group whose order are power of  $p$ . This number  $r_p$  is the  $p$ -rank of  $p$ -subgroup of  $C$ . Observe that if

$p^{e_p} \parallel h(L)$  then  $r_p \leq e_p$  and that  $r_p$  is generally *small* and more often  $r_p = 1$ .

## 1.2 Some results

**Theorem 1.1.** *Let  $L/\mathbb{Q}$  be a finite algebraic extension with  $[L : \mathbb{Q}] = 2^{\alpha_0} \times N_1$ , where  $N_1 > 1$  is odd. Suppose that there exists a field  $K \subset L$  with  $[K : \mathbb{Q}] = 2^{\alpha_0}$  and with  $L/K$  Galois solvable extension. Let  $h(L)$  be the class number of  $L$ . Suppose that  $h(L) > 1$ . Let  $p$  be a prime dividing  $h(L)$ . Let  $r_p$  be the rank of the  $p$ -class group of  $L$ . If  $p \times \prod_{i=1}^{r_p} (p^i - 1)$  and  $N_1$  are coprime, then  $p$  divides the class number  $h(K)$  of  $K$ .*

*Proof.*

- $L/K$  being Galois solvable, we can always suppose that there exists a tower of fields

$$K \subset L_1 \subset L_2 \subset \cdots \subset L_j \subset L_{j+1} \subset \cdots \subset L_t \subset L,$$

such that  $[L_{j+1} : L_j] = q_j$  where  $L_{j+1}/L_j$ ,  $j = 1, \dots, t$ , is a cyclic extension with  $q_j$  prime.

- The extension  $L/L_t$  is cyclic with  $[L : L_t] = q_t$  and  $q_t | N_1$ .
- Suppose at first that  $p \nmid h(L_t)$ : then hypotheses of rank theorem are verified, see Masley [8] corollary 2.15 p. 305 :
  - From hypothesis,  $p$  does not divide  $N_1$ , so  $p$  does not divide  $q_t = [L : L_t]$ .
  - The extension  $L/L_t$  is cyclic with  $h(L) \equiv 0 \pmod{p}$ ,  $h(L_t) \not\equiv 0 \pmod{p}$ .
  - There is no field  $E$ , different of  $L$  and of  $L_t$  with  $L_t \subset E \subset L$ .

From rank theorem, if  $f$  is the order of  $p \pmod{q_t}$  then  $f | r_p$  and so  $p^{r_p} - 1 \equiv 0 \pmod{q_t}$ , which contradicts hypothesis.

- Therefore  $p$  divides  $h(L_t)$  : the  $p$ -rank  $r'_p$  of class group  $h(L_t)$  of  $L_t$  verifies  $r'_p \leq r_p$  because  $h(L_t) \mid q_t \times h(L)$ , see for instance Masley, [8], *pushing up* corollary 2.2 p. 301 and so  $p$  and  $q_t$  coprime implies that the  $p$ -class group  $C_p(L_t)$  of  $L_t$  and  $C_p$  of

$L$  verify  $C_p(L_t) \subset C_p(L)$  and so  $r'_p \leq r_p$ . Then let us consider  $N' = \frac{N}{q_t} = [L_t : K]$ . We can pursue the same algorithm with same Masley corollaries 2.15 and 2.2 applied to extension  $L_t/K$  in place of  $L/K$  and  $N'$  in place of  $N$  and with  $r'_p \leq r_p$  in place of  $r_p$ , up to find a prime divisor  $q_j$ ,  $1 \leq j \leq t$ , of  $N_1$  dividing  $\prod_{i=1}^{r_p} (p^i - 1)$ , or to get the subfield  $K$  of  $L$  with  $[K : \mathbb{Q}] = 2^{\alpha_0}$  and  $p \mid h(K)$ , which achieves the proof.

□

In following corollary, we generalize the result obtained for  $L/K$  Galois solvable extension to the case where  $L/K$  is a solvable extension.

**Corollary 1.2.** *Let  $L/\mathbb{Q}$  be a finite algebraic extension with  $[L : \mathbb{Q}] = 2^{\alpha_0} \times N_1$ , where  $N_1 > 1$  is odd. Suppose that there exists a field  $K \subset L$  with  $[K : \mathbb{Q}] = 2^{\alpha_0}$  and with  $L/K$  solvable extension. Let  $M/K$  be the Galois solvable extension containing  $L/K$ . Let  $h(L)$  be the class number of  $L$ . Suppose that  $h(L) > 1$ . Let  $p$  be a prime dividing  $h(L)$  such that  $p > [L : K]$ . Let  $r_p(M)$  be the rank of the  $p$ -class group of  $M$ . If  $\prod_{i=1}^{r_p(M)} (p^i - 1)$  and  $N_1$  are coprime, then  $p$  divides the class number  $h(K)$  of  $K$ .*

*Proof.* From *pushing up* theorem, see Masley [8] corollary 2.2 p. 301,  $h(L) \mid [M : L] \times h(M)$ . Let  $q_M$  be the greatest prime divisor of  $[M : K]$ . The solvable extension  $M/K$  is the smallest Galois extension of  $K$  containing  $L$ . Therefore  $q_M \mid [L : K]!$ , see for instance Morandi, [9], cor. 3.8 p. 29. From hypothesis, this implies that  $p \nmid [M : L]$ . Therefore  $p \mid h(M)$ . Then apply theorem 1.1 p. 5 to Galois solvable extension  $M/K$ . □

In following corollary we give a particularly straightforward formulation of theorem 1.1 p. 5 when  $[L : \mathbb{Q}] = N$  is odd.

**Corollary 1.3.** *Let  $L/\mathbb{Q}$  be a Galois solvable extension with  $[L : \mathbb{Q}] = N$ , where  $N > 1$  is odd. Let  $h(L)$  be the class number of  $L$ . Suppose that  $h(L) > 1$ . Let  $p$  be a prime dividing  $h(L)$ . Let  $r_p$  be the rank of the  $p$ -class group of  $L$ . Then  $p \times \prod_{i=1}^{r_p} (p^i - 1)$  and  $N$  are not coprime.*

*Proof.* Apply theorem 1.1 p. 5. □

**Remark:** Observe that, with previous notations, if  $L/K$  is abelian then  $L/K$  is Galois solvable and so the previous results can be applied to all abelian extensions  $L/K$  with  $[K : \mathbb{Q}] = 2^{\alpha_0}$  and  $[L : K]$  odd.

## 2 Geometry of Numbers point of view

- The next result connects rank theorem to Geometry of Numbers point of view. Let  $L/\mathbb{Q}$  be a finite algebraic extension. Suppose that there exists a subfield  $F$  of  $L$  such that  $L/F$  is a cyclic extension, with  $[L : \mathbb{Q}] = N$ ,  $[L : F] = q$  where  $q$  is an odd prime.
- Let  $D_F$  be the discriminant of  $F$ . Let  $m = \frac{N}{q}$ . Then  $h(F) \leq \frac{2^{m-1}}{(m-1)!} \times \sqrt{|D_F|} \times (\log(|D_F|))^{m-1}$ , see Bordellès [2] theorem 5.3 p. 4. Let us note

$$(1) \quad H_F = \frac{2^{m-1}}{(m-1)!} \times \sqrt{|D_F|} \times (\log(|D_F|))^{m-1}.$$

If a prime  $p$  verifies  $p > H_F$  then  $p \nmid h(F)$ .

**Theorem 2.1.** *Let  $L/\mathbb{Q}$  be a finite algebraic extension. Let  $h(L)$  be the class number of  $L$ . Suppose that  $h(L) > 1$ . Suppose that there exists a cyclic extension  $L/F$ , where  $q = [L : F]$  is an odd prime. Let  $H_F$  be a geometric upper bound of class number of  $F$  given by relation (1). Suppose that  $p > H_F$  is a prime dividing  $h(L)$ . Let  $r_p$  be the rank of the  $p$ -class group of  $L$ . Then  $p \times (p^{r_p} - 1) \equiv 0 \pmod{q}$ .*

*Proof.* We apply rank theorem, see Masley[8] corollary 2.15 p 305, and upper bound  $H_F$  of class number  $h(F)$  of field  $F$  given in relation (1).  $\square$

**Corollary 2.2.** *Let  $L/\mathbb{Q}$  be a finite Galois extension with  $[L : \mathbb{Q}] = N$ . Let  $h(L)$  be the class number of  $L$ . Suppose that  $h(L) > 1$ . Suppose that  $N$  has odd prime divisors. Let  $q$  be an odd prime divisor of  $N$ . Then there exists a cyclic extension  $L/F$  with  $q = [L : F]$ . Let  $H_F$  be a geometric upper bound of class number of  $F$  given by relation (1). Suppose that  $p > H_F$  is a prime dividing  $h(L)$ . Let  $r_p$  be the rank of the  $p$ -class group of  $L$ . Then  $p \times (p^{r_p} - 1) \equiv 0 \pmod{q}$ .*

*Proof.* Immediate consequence of theorem 2.1 and of Galois theory.  $\square$

**Remark:** Observe that all the results above give strong information on prime factor of class numbers of Galois solvable extensions of  $L/K$ , because practically the  $p$ -rank  $r_p$  is generally *small*: in all numerical examples following  $r_p \leq 3$  and it verifies more often  $r_p = 1$ .

### 3 Numerical examples

The examples found to check theses results are taken from:

- the table of relative class numbers of cyclotomic number fields in Washington, [11] p 412, with some elementary MAPLE computations,
- the table of relative class number of cyclotomic number fields in Schoof, [10]
- the table of maximal real subfields  $\mathbb{Q}(\zeta_l + \zeta_l^{-1})$  of  $\mathbb{Q}(\zeta_l)$  for  $l$  prime in Washington, [11] p 420.
- the tables of relative class number of imaginary cyclic fields of Girstmair of degree 4,6,8,10. [4].
- the tables of quintic number fields computed by Jeannin, [5]
- the tables of cubic totally real cyclic number fields of the Bordeaux University in the Server  
*megrez.math.u-bordeaux.fr*.

All the results examined in these tables are in accordance with our theorems.

#### 3.1 Cyclotomic number fields $\mathbb{Q}(\zeta_u)$

Let  $u \in \mathbb{N}$ ,  $u > 2$ . Let  $\zeta_u$  be a primitive  $u^{th}$  root of unity. Here we have  $N = \phi(u)$ , where  $\phi$  is the Euler indicator. The cyclotomic number fields  $\mathbb{Q}(\zeta_u)$  of the examples are taken with  $2 \parallel N = \phi(u)$ , except the example  $\zeta_u$  with  $u = 572$ ,  $N = \phi(u) = 2^4.3.5$ .

- $\mathbb{Q}(\zeta_u)$ ,  $u = 59$ ,  $N = \phi(u) = 58 = 2.29$  :  
 $h^- = 3.(2.29 + 1).(2^3.29 + 1)$ ,  $h^+ = 1$ .  
 $3 = h(\mathbb{Q}(\sqrt{-59}))$ .



- $\mathbb{Q}(\zeta_u)$ ,  $u = 71$ ,  $N = \phi(u) = 70 = 2.5.7$  :  
 $h^- = 7^2.(2^3.5.7.283 + 1)$ ,  $h^+ = 1$ .
- $\mathbb{Q}(\zeta_u)$ ,  $u = 79$ ,  $N = \phi(u) = 78 = 2.3.13$  :  
 $h^- = 5.(2^2.13 + 1)(2.3^2.5.13.17.19 + 1)$ ,  $h^+ = 1$ .  
 $5 = h(\mathbb{Q}(\sqrt{-79}))$ .
- $\mathbb{Q}(\zeta_u)$ ,  $u = 83$ ,  $N = \phi(u) = 82 = 2.41$  :  
 $h^- = 3.(2^2.41.1703693 + 1)$ ,  $h^+ = 1$ .  
 $3 = h(\mathbb{Q}(\sqrt{-83}))$ .
- $\mathbb{Q}(\zeta_u)$ ,  $u = 103$ ,  $N = \phi(u) = 102 = 2.3.17$ :  
 $h^- = 5.(2.3.17 + 1)(2^2.3.5.17 + 1)(2.3^2.5.17.11273 + 1)$ ,  
 $h^+ = 1$ .
- $\mathbb{Q}(\zeta_u)$ ,  $u = 131$ ,  $N = \phi(u) = 130 = 2.5.13$  :  
 $h^- = 3^3.5^2.(2^2.13 + 1)(2.5.13 + 1)(2^2.5^2.13 + 1) \times$   
 $(2^2.5^2.13.29.151.821 + 1)$ ,  $h^+ = 1$ .

Observe that for  $p = 3$ , we have  $p^3 \parallel h$  and the group  $C_p$  is not cyclic as it is seen in Schoof [10] table 4.2 p 1239, where  $r_p = 3$ . The rank theorem in that case shows that  $f = r_p = 3$  and so that  $3^f = 27 \equiv 1 \pmod{13}$ .

- $\mathbb{Q}(\zeta_u)$ ,  $u = 151$ ,  $N = \phi(u) = 150 = 2.3.5^2$  :  
 $h^- = (2.3 + 1)(2.5 + 1)^2.(2^2.5.7 + 1)(2.3.5^2.173 + 1) \times$   
 $(2^2.3.5^4.7.23 + 1)(2^2.3.5^2.7.13.73.1571 + 1)$ ,  $h^+ = 1$ .

For  $p = 2.5 + 1 = 11$ , from Schoof, see table 4.2 p 1239, the group is not cyclic,  $r_p = 2$ . From rank theorem, we can only assert that  $f | r_p = 2$ , so that  $f = 1$  or  $f = 2$ , so only that  $p^2 \equiv 1 \pmod{g = 5}$ . We see in this numerical application that in that case  $f = 1$  and  $f < r_p$  strictly.

- $\mathbb{Q}(\zeta_u)$ ,  $u = 191$ ,  $N = \phi(u) = 190 = 2.5.19$ :  
 $h^- = (2.5 + 1).13.(2.19^2.71 + 1)(2.3.5.19.277.3881 + 1)$ ,  
 $h^+ = (2.5 + 1)$ .  
 $13 = h(\mathbb{Q}(\sqrt{-191}))$ .

Here, we note that the prime  $p = 11$  corresponds to 11-class group of  $\mathbb{Q}(\zeta_{191} + \zeta_{191}^{-1})$ .

- $\mathbb{Q}(\zeta_u)$ ,  $u = 572 = 2^2.11.13$ ,  $N = \phi(u) = 2^4.3.5$ ,  
 $h^- = 3.5^2.(2.3 + 1)(2.3^2 + 1)^2(2.3.5 + 1)(2^3.5 + 1)(2^2.3.5 + 1)^2$   
 $(2^2.3.5.7 + 1)(2^2.3.5.11 + 1)(2.3^2.5.307 + 1)(2.3^2.5.11.73 + 1)$   
 $(2^2.3^2.5^2.31 + 1)(2^2.3^5.5.7.53.263 + 1)$

We don't know  $h^+$ .

#### Remarks:

- Let  $[K : \mathbb{Q}] = 2^{\alpha_0} \times n_1^{\alpha_1} \times \cdots \times n_k^{\alpha_k}$ . We observe in Washington [11], tables of relative class numbers p. 412 and of real class numbers p. 421 that, when  $p$  is large,  $p-1$  is divisible by several or all primes in the set  $\{n_i \mid i = 1, \dots, k\}$ . This observation complies with Geometry of Number corollary 2.2 p. 7.
- Observe that frequently, we get  $f = r_p = 1$  and so  $p \times (p-1) \equiv 1 \pmod n$ . In our examples we get only one example  $\mathbb{Q}(\zeta_{131})$  with  $p = 3$ ,  $f = r_p = 3$ .

### 3.2 Real class number $\mathbb{Q}(\zeta_l + \zeta_l^{-1})$

The examples are obtained from the table of real class number in Washington, [11] p 420. Here,  $l$  is a prime, the class number  $h_\delta$  is the conjectured value of the class number  $h^+$  of  $\mathbb{Q}(\zeta_l + \zeta_l^{-1})/\mathbb{Q}$  with a minor incertitude on an extra factor. But the factor  $h_\delta$  must verify our theorems. We extract some examples of the table with  $2 \parallel l-1$ . Here we have  $N = \frac{l-1}{2}$  and thus  $N \not\equiv 0 \pmod 2$ .

- $l = 191$ ,  $l-1 = 2.5.19$ ,  $h_\delta = (2.5 + 1)$
- $l = 1063$ ,  $l-1 = 2.3^2.59$ ,  $h_\delta = (2.3 + 1)$ .
- $l = 1231$ ,  $l-1 = 2.3.5.41$ ,  $h_\delta = (2.3.5.7 + 1)$ .
- $l = 1459$ ,  $l-1 = 2.3^6$ ,  $h_\delta = (2.3.41 + 1)$ .
- $l = 1567$ ,  $l-1 = 2.3^3.29$ ,  $h_\delta = (2.3 + 1)$ .
- $l = 2659$ ,  $l-1 = 2.3.443$ ,  $h_\delta = (2.3^2 + 1)$ .
- $l = 3547$ ,  $l-1 = 2.3^2.197$ ,  $h_\delta = ((2.3^2 + 1)(2.3^2.7^2 + 1))$ .
- $l = 8017$ ,  $l-1 = 2^4.3.167$ ,  
 $h_\delta = (2.3^2 + 1)(2.3^2.7^2 + 1)(2^2.3^3 + 1)$ .

- $l = 8563$ ,  $l - 1 = 2.3.1427$ ,  $h_\delta = (2.3 + 1)^2$ . We have  $r_p = 1$  or  $r_p = 2$  so  $f = 1$  or  $f = 2$ . We can conclude from theorem 1.1 p. 5, that  $p^2 \equiv 1 \pmod{3}$ .
- $l = 9907$ ,  $l - 1 = 2.3.13.127$ ,  $h_\delta = (2.3.5 + 1)$ .

### 3.3 Cubic fields $K/\mathbb{Q}$ cyclic and totally real.

Here, we have  $N = 3$ . Note that in that case discriminants are square in  $\mathbb{N}$ .

- Discriminant  $D_1 = 3969 = (3^2.7)^2$ ,  $h = 3$
- Discriminant  $D_2 = 3969 = (3^2.7)^2$ ,  $h = 7 = 2.3 + 1$
- Discriminant  $D_1 = 8281 = (7.13)^2$ ,  $h = 3$

### 3.4 Totally real cyclic fields of prime conductor $< 100$

We have found few numeric results in the literature. We refer to Masley, [8], Table 3 p 316. In these results,  $K/\mathbb{Q}$  is a real cyclic field with  $[K : \mathbb{Q}] = N$ , with conductor  $f$ , with root of discriminant  $Rd$  and class number  $h$ .

- $f = 63$ ,  $N = 3$ ,  $Rd = 15.84$ ,  $h = 3$
- $f = 63$ ,  $N = 3$ ,  $Rd = 15.84$ ,  $h = 3$
- $f = 63$ ,  $N = 6$ ,  $Rd = 26.30$ ,  $h = 3$
- $f = 63$ ,  $N = 6$ ,  $Rd = 26.30$ ,  $h = 3$
- $f = 91$ ,  $N = 3$ ,  $Rd = 20.24$ ,  $h = 3$

### 3.5 Lehmer quintic cyclic field

The prime divisors of the 82 cyclic number fields of the table in Jeannin [5], with conductor  $f < 3000000$ , are, at a glance, of the form  $p = 2$  or  $p = 5$  or  $p \equiv 1 \pmod{10}$ , which clearly verifies corollary 1.3 p.6.

### 3.6 Decimic imaginary cyclic number fields with conductor between 9000 and 9500

This example is obtained from the tables of Girstmair, [4].  $f$  is a prime conductor,  $h$  is the factorization of the class number  $K/\mathbb{Q}$ .

- $f = 9011$ ,  $h = 3.(2.5 + 1).(2.3.5.52201 + 1)$ .  
3 divides the class number of  $\mathbb{Q}(\sqrt{-9011})$ .
- $f = 9081$ ,  $h = 3.7.(2^3.5 + 1)$ .  
3, 7 divide the class number of  $\mathbb{Q}(\sqrt{-9081})$ .
- $f = 9151$ ,  $h = 67.(2^2.5.1187 + 1)$ .  
67 divides the class number of  $\mathbb{Q}(\sqrt{-9151})$ .
- $f = 9311$ ,  $h = 97.(2.5.5689 + 1)$ .  
97 divides the class number of  $\mathbb{Q}(\sqrt{-9311})$ .
- $f = 9371$ ,  $h = 7^2.(2.3.5^2 + 1).(2.3^3.5 + 1)$ .  
7 divides the class number of  $\mathbb{Q}(\sqrt{-9371})$ .
- $f = 9391$ ,  $h = 5^2.(2^4.3.5.7.71 + 1)$ .

### 3.7 Not abelian Galois solvable extensions $L/K$

Here, we give an example of Galois solvable extension not abelian. Let  $q$  be a prime with  $q = 2g + 1$ , where  $g$  is an odd prime. Let  $\zeta_q$  be a root of the equation  $X^{q-1} + \cdots + X + 1 = 0$  and let  $\omega$  root of the equation  $X^q - 2 = 0$ . The extension  $L = \mathbb{Q}(\zeta_q, \omega)/\mathbb{Q} = \mathbb{Q}(\zeta_q)(\omega)$  verifies  $\mathbb{Q} \subset \mathbb{Q}(\sqrt{-q}) \subset \mathbb{Q}(\zeta_q) \subset L$ , where  $\mathbb{Q}(\zeta_q)/\mathbb{Q}(\sqrt{-q})$  is cyclic and where  $\mathbb{Q}(\zeta_q, \omega)/\mathbb{Q}(\zeta_q)$  is cyclic. Let  $p$  be a prime dividing the class number  $h(L)$  with  $p \neq g$ ,  $p \neq q$ . Apply theorem 1.1 p. 5 to get:

Let  $r_p$  be the rank of the  $p$ -class group of  $L$ . Then else  $p \mid h(\mathbb{Q}(\sqrt{-q}))$  else  $Gcd(\prod_{i=1}^{r_p} (p^i - 1), q \times g) > 1$ .

## References

- [1] J.L. Alperin, Rowen B. Bell, *Groups and Representations*, Springer, 1995.

- [2] O. Bordellès, *Explicit upper bounds for the average order of  $d_n(m)$  and application to class number*, Journal of Inequalities in Pure and Applied Mathematics, vol 3, issue 3, article 38, 2002, <http://jipam.vu.edu.au/>
- [3] J. Calais, *éléments de théorie des groupes*, Presses Universitaires de France, 1984.
- [4] K. Girstmair, *The relative class number of imaginary cyclic fields of degree 4, 6, 8 and 10*, Math. Comp., 61, 204, 1993, p 881-887.
- [5] S. Jeannin, *Nombre de classes et unités des corps de nombres cycliques quintiques d'E. Lehmer*, J. Théorie des Nombres de Bordeaux, 1996, 8, pp75-92.
- [6] S. Lang, *Algebra*, Addison-Wesley, 1978.
- [7] D.H. Lehmer, *Prime factors of cyclotomic class numbers*, Math. Comp., 31, 138, 1977, pp 599-607.
- [8] J.M. Masley, *Class number of real cyclic number fields with small conductor*, Compositio Mathematica, 37, 3., 1978, pp 297-319.
- [9] P. Morandi, *Field and Galois Theory*, Springer, 1996.
- [10] R. Schoof, *Minus class groups of the fields of  $l$ -th roots of unity*, Math. Comp., 67, 223, 1998, pp 1225-1245.
- [11] L.C. Washington, *Introduction to cyclotomic field, second edition*, Springer, 1997.

\*\*\*\*\*

Roland Quême  
 13 avenue du château d'eau  
 31490 Brax  
 France  
 2003 avril 20  
 tel 0561067020  
 e-mail : roland.queme@free.fr  
 home page: <http://roland.queme.free.fr/>